FILDES LABELS

Noroxin Label 3b, 4, 8, 12	3b Take on an **empty stomach** at least half an hour before food or two hours after food. 4 Do not take dairy products, antacids, iron or calcium supplements within two hours of each dose of this medicine. 8 Avoid excessive skin exposure to sunlight and sunlamps while being treated with this medicine. 12 This medicine may affect mental alertness and/or coordination. If affected, do not drive a motor vehicle or operate machinery.
Doxy Label 4 (delete diary products), 8, B	4 Do not take ~~dairy products~~, antacids, iron or calcium supplements within two hours of each dose of this medicine. B - Take with or soon after food.
Temazepam Label 1 or 1a, 9	1 THIS MEDICINE MAY CAUSE DROWSINESS AND MAY INCREASE THE EFFECTS OF ALCOHOL. IF AFFECTED DO NOT DRIVE A MOTOR VEHICLE OR OPERATE MACHINERY. 9 DO NOT STOP TAKING THIS MEDICINE ABRUPTLY unless otherwise advised by your doctor.
Simplotan Label 2, B	2 **DO NOT TAKE ALCOHOL** while being treated with this medicine. B - Take with or soon after food.
Stemzine Label 1	1 THIS MEDICINE MAY CAUSE DROWSINESS AND MAY INCREASE THE EFFECTS OF ALCOHOL. IF AFFECTED DO NOT DRIVE A MOTOR VEHICLE OR OPERATE MACHINERY.
Stilnox Label 1 or 1a, 9, 12	1 THIS MEDICINE MAY CAUSE DROWSINESS AND MAY INCREASE THE EFFECTS OF ALCOHOL. IF AFFECTED DO NOT DRIVE A MOTOR VEHICLE OR OPERATE MACHINERY. 9 DO NOT STOP TAKING THIS MEDICINE ABRUPTLY unless otherwise advised by your doctor. 12 This medicine may affect mental alertness and/or coordination. If affected, do not drive a motor vehicle or operate machinery.
Rulide Label 3b	3b Take on an **empty stomach** at least half an hour before food or two hours after food.
Bactrim Label 8, B	8 Avoid excessive skin exposure to sunlight and sunlamps while being treated with this medicine.
Augmentin Due Forte Label F	Immediately before food.
Diamox Label 10a	Do not take more than 1 aspirin tablet/capsule each day while being treated with this medicine

B - Take with or soon after food.